IN THE CLAIMS

Please amend claim 21 as follows.

This listing of the claims replaces all prior versions of the claims in the application.

1-20. (Canceled)

- 21. (Currently Amended) An isolated polypeptide selected from the group consisting of:
- a) a polypeptide comprising an amino acid sequence of SEQ ID NO:17,
- a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical to an amino acid sequence of SEQ ID NO:17,
- e) a biologically active fragment of a polypeptide having an amino acid sequence of SEQ ID NO:17, and
- d) c) an immunogenic fragment of a polypeptide consisting of an amino acid sequence of SEQ ID NO:17, wherein said fragment comprises at least 40 contiguous amino acid residues of SEQ ID NO:17.
- 22. (Previously Presented) An isolated polynucleotide encoding a polypeptide of claim 21.
- 23. (Previously Presented) An isolated polynucleotide of claim 22 comprising a polynucleotide sequence of SEQ ID NO:44.
- 24. (Previously Presented) A recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide of claim 22.
- 25. (Previously Presented) A cell transformed with a recombinant polynucleotide of claim 24.
- 26. (Previously Presented) A method of producing a polypeptide of claim 21, the method comprising:

120125 2 09/857,826

- a) culturing a cell under conditions suitable for expression of the polypeptide, wherein said cell is transformed with a recombinant polynucleotide, and said recombinant polynucleotide comprises a promoter sequence operably linked to a polynucleotide encoding the polypeptide of claim 21, and
- b) recovering the polypeptide so expressed.
- 27. (Previously Presented) A method of claim 26, wherein the polypeptide comprises an amino acid sequence of SEQ ID NO:17.
- 28. (Previously Presented) An isolated polynucleotide selected from the group consisting of:
 - a) a polynucleotide comprising a polynucleotide sequence of SEQ ID NO:44,
 - b) a polynucleotide comprising a naturally occurring polynucleotide sequence at least 90% identical to a polynucleotide sequence of SEQ ID NO:44,
 - c) a polynucleotide complementary to a polynucleotide of a),
 - d) a polynucleotide complementary to a polynucleotide of b), and
 - e) an RNA equivalent of a)-d).
- 29. (Withdrawn) A method of detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide of claim 28, the method comprising:
 - a) hybridizing the sample with a probe comprising at least 20 contiguous nucleotides comprising a sequence complementary to said target polynucleotide in the sample, and which probe specifically hybridizes to said target polynucleotide, under conditions whereby a hybridization complex is formed between said probe and said target polynucleotide or fragments thereof, and
 - b) detecting the presence or absence of said hybridization complex, and, optionally, if present, the amount thereof.
- 30. (Withdrawn) A method of detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide of claim 28, the method comprising:

120125 3 09/857,826

- a) amplifying said target polynucleotide or fragment thereof using polymerase chain reaction amplification, and
- b) detecting the presence or absence of said amplified target polynucleotide or fragment thereof, and, optionally, if present, the amount thereof.
- 31. (Previously Presented) A composition comprising a polypeptide of claim 21 and a pharmaceutically acceptable excipient.
- 32. (Previously Presented) A composition of claim 31, wherein the polypeptide comprises an amino acid sequence of SEQ ID NO:17.
- 33. (Withdrawn) A method of screening for a compound that specifically binds to the polypeptide of claim 21, the method comprising:
 - a) combining the polypeptide of claim 21 with at least one test compound under suitable conditions, and
 - b) detecting binding of the polypeptide of claim 21 to the test compound, thereby identifying a compound that specifically binds to the polypeptide of claim 21.
- 34. (Withdrawn) A method of screening a compound for effectiveness in altering expression of a target polynucleotide, wherein said target polynucleotide comprises a sequence of claim 23, the method comprising:
 - a) exposing a sample comprising the target polynucleotide to a compound, under conditions suitable for the expression of the target polynucleotide,
 - b) detecting altered expression of the target polynucleotide, and
 - c) comparing the expression of the target polynucleotide in the presence of varying amounts of the compound and in the absence of the compound.
- 35. (Withdrawn) A method of assessing toxicity of a test compound, the method comprising:
 - a) treating a biological sample containing nucleic acids with the test compound,

120125 4 09/857,826

- b) hybridizing the nucleic acids of the treated biological sample with a probe comprising at least 20 contiguous nucleotides of a polynucleotide of claim 28 under conditions whereby a specific hybridization complex is formed between said probe and a target polynucleotide in the biological sample, said target polynucleotide comprising a polynucleotide sequence of a polynucleotide of claim 28 or fragment thereof,
- c) quantifying the amount of hybridization complex, and
- d) comparing the amount of hybridization complex in the treated biological sample with the amount of hybridization complex in an untreated biological sample, wherein a difference in the amount of hybridization complex in the treated biological sample is indicative of toxicity of the test compound.
- 36. (Withdrawn) A microarray wherein at least one element of the microarray is a polynucleotide of claim 22.
- 37. (Withdrawn) A method of generating an expression profile of a sample which contains polynucleotides, the method comprising:
 - a) ' labeling the polynucleotides of the sample,
 - contacting the elements of the microarray of claim 36 with the labeled
 polynucleotides of the sample under conditions suitable for the formation of a
 hybridization complex, and
 - c) quantifying the expression of the polynucleotides in the sample.
- 38. (Withdrawn) An isolated antibody which specifically binds to a polypeptide of claim 21.
 - 39. (Withdrawn) The antibody of claim 38, wherein the antibody is:
 - a) a chimeric antibody,
 - b) a single chain antibody,
 - c) a Fab fragment,
 - d) a F(ab')₂ fragment, or

e) a humanized antibody.

40. (Withdrawn) A composition comprising an antibody of claim 38 and an acceptable excipient.